

Xylanase 10C, *Clostridium thermocellum*

CtXyn10C (CBM22-GH10)

Catalogue number	Presentation
CZ00051	1 mg
CZ00052	3 x 1 mg

Description

Xylanase 10C (CtXyn10C), assigned the E.C. number 3.2.1.8, is a derivative of *Clostridium thermocellum*. It is an endo-1,4- β -xylanase. The recombinant CtXyn10C, purified from *Escherichia coli*, is a modular Glycoside Hydrolase family 10 (CBM22-GH10) enzyme (see more details at www.cazy.org). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl₂, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

Electrophoretic Purity

The molecular integrity and purity of CtXyn10C (CBM22-GH10) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).




Figure 1. SDS-PAGE analysis of CtXyn10C (CBM22-GH10) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 59,32 kDa. Lane M contains a Protein Marker for reference.

Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

Substrate specificity

CtXyn10C (CBM22-GH10) hydrolyses xylan, such as arabinoxylan and oat spelt xylan.

Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 4.0-11.0 and at a temperature of 80°C. Maximal enzymatic activity is achieved at pH 5.5 and a consistent temperature of 80°C.

Specific activity

CtXyn10C (CBM22-GH10) specific activity is 450 U/mg, using wheat arabinoxylan as substrate.

Enzyme activity

Substrate specificity and kinetic properties of CtXyn10C (CBM22-GH10) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

Reference

Hayashi *et al.* (1997) J. Bacteriol. 179, 4246-4253.

Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

Quality control assay

Protein purity is determined to be $\geq 90\%$, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.