

Exo-1,3- β -Galactanase 43A, *Clostridium thermocellum*

CtGan43A (GH43)

Catalogue number	Presentation
CZ00291	2 mg
CZ00292	3 x 2 mg

Description

Exo-1,3- β -Galactanase 43A (CtGan43A), assigned the E.C. number 3.2.1.145, is a derivative of *Clostridium thermocellum*. It is an exo- α -arabinofuranosidase. The recombinant CtGan43A, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 43 (GH43) enzyme (see more details at www.cazy.org). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl₂, and 25% (v/v) glycerol, at a concentration of 2 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

Electrophoretic Purity

The molecular integrity and purity of CtGan43A (GH43) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).




Figure 1. SDS-PAGE analysis of CtGan43A (GH43) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 37,85 kDa. Lane M contains a Protein Marker for reference.

Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

Substrate specificity

CtGan43A (GH43) hydrolyses 1,3- β -galactans in an exo-acting manner.

Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 3.0-10.0 and at a temperature of 50°C. Maximal enzymatic activity is achieved at pH 6 and a consistent temperature of 50°C.

Specific activity

CtGan43A (GH43) specific activity is 20 U/mg, using β -1,3-galactan as substrate.

Enzyme activity

Substrate specificity and kinetic properties of CtGan43A (GH43) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

Reference

Ichinose *et al.* (2006) *Appl. Envir. Microbiol.* 72, 3515-3523.

Jiang *et al.* (2012) *J.Struct.Biol.* 180: 447-457.

Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

Quality control assay

Protein purity is determined to be $\geq 90\%$, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.

NZYtech Lda. Estrada do Paço do Lumiar, Campus do Lumiar - Edifício E, R/C, 1649-038 Lisboa, Portugal Tel.:+351.213643514 Fax:
+351.217151168 www.nzytech.com