

CZ0111_UG_EN_V2302

Oligo-1,6-α-Glucosidase 13B, Bacillus subtilis

BsAgl13B (GH13)

 Catalogue number
 Presentation

 CZ01111
 0.25 mg

 CZ01112
 3 x 0.25 mg

Description

Oligo-1,6- α -Glucosidase 13B (BsAgl13B), assigned the E.C. number 3.2.1.10, is a derivative of $Bacillus\ subtilis$. It is an oligosaccharide 1,6- α -glucosidase. The recombinant BsAgl13B, purified from $Escherichia\ coli$, is a single-domain Glycoside Hydrolase family 13 (GH13) enzyme (see more details at www.cazy.org). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl₂, and 25% (v/v) glycerol, at a concentration of 0.25 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

Electrophoretic Purity

The molecular integrity and purity of BsAgl13B (GH13) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).




Figure 1. SDS-PAGE analysis of *Bs*Agl13B (GH13) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 68,11 kDa. Lane M contains a Protein Marker for reference.

Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

Substrate specificity

BsAgl13B (GH13) hydrolyses 1,6- α -glucosides.

Temperature and pH optima

The pH optimum for enzymatic activity is 7 while temperature optimum is 25 °C.

Enzyme activity

The substrate specificity and kinetic properties of *Bs*Agl13B (GH13) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

Reference

Schönert et al. (1998) J Bacteriol. 180(9):2574-2578.

Schönert et al. (1999) Res Microbiol. 150(3):167-177.

Hobbs et al. (2013) ACS Chem Biol. 8(11):2388-93.

Kamp et al. (2018) Nat Commun. 9(1):1177.

Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

Quality control assay

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.