

CZ0129\_UG\_EN\_V2302

# Unsaturated rhamnogalacturonyl hydrolase 105A, Bacillus subtilis

# BsUrh105A (GH105)

Catalogue number Presentation


CZ01291 1 mg CZ01292 3 x 1 mg

#### **Description**

Unsaturated rhamnogalacturonyl hydrolase 105A (*Bs*Urh105A), assigned the E.C. number 3.2.1.172, is a derivative of *Bacillus subtilis*. It catalyzes the hydrolysis of unsaturated rhamnogalacturonan disaccharide to yield unsaturated D-galacturonic acid and L-rhamnose. The recombinant *Bs*Urh105A, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 105 (GH105) enzyme (see more details at <a href="https://www.cazy.org">www.cazy.org</a>). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

## **Electrophoretic Purity**

The molecular integrity and purity of BsUrh105A (GH105) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1**. SDS-PAGE analysis of *Bs*Urh105A (GH105) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 43,48 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### **Substrate specificity**

BsUrh105A (GH105) hydrolyses unsaturated rhamnogalacturonan (RG).

# Temperature and pH optima

The pH optimum for enzymatic activity is 4 while temperature optimum is 30 °C.

# **Enzyme activity**

The substrate specificity and kinetic properties of BsUrh105A (GH105) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

#### Reference

Itoh et al. (2006a) J. Mol. Biol. 360:573-585.

Itoh et al. (2006b) Biochem Biophys Res Commun. 347(4):1021-1029.

#### **Customer Support**

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

## **Quality control assay**

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.