

CZ0145_UG_EN_V2302

α-Mannosidase 38A, Bacteroides thetaiotaomicron

BtMns38A (GH38)

Catalogue number Presentation

CZ01451 1 mg CZ01452 3 x 1 mg

Description

 α -Mannosidase 38A (BtMns38A), assigned the E.C. number 3.2.1.113 and 3.2.1.-, is a derivative of Bacteroides thetaiotaomicron. It is an unsepecific mannosyl α -mannosidase acting 1,2 and 1,3 and 1,6 linkages of α -mannans. The recombinant BtMns38A, purified from Escherichia coli, is a single-domain Glycoside Hydrolase family 38 (GH38) enzyme (see more details at www.cazy.org). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl₂, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

Electrophoretic Purity

The molecular integrity and purity of BtMns38A (GH38) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).




Figure 1. SDS-PAGE analysis of *Bt*Mns38A (GH38) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 134,3 kDa. Lane M contains a Protein Marker for reference.

Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

Substrate specificity

BtMns38A (GH38) hydrolyses yeast α-mannans.

Temperature and pH optima

The pH optimum for enzymatic activity is 7.5 while temperature optimum is 37 °C.

Enzyme activity

The substrate specificity and kinetic properties of *Bt*Mns38A (GH38) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

Reference

Cuskin et al. (2015) Nature 517, 165-169.

Mahowald et al. (2009) Proc Natl Acad Sci U S A. 106(14):5859-5864.

Nankai et al. (2002) Appl Environ Microbiol. 68(6):2731-2736.

Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

Quality control assay

Protein purity is determined to be ≥75%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.