

CZ0212 UG EN V2302

# Phospho-β-Glucosidase 4A, Thermotoga maritima

# TmPbg4A (GH4)

Catalogue number Presentation


CZ02121 1 mg CZ02122 3 x 1 mg

### **Description**

Phospho-β-Glucosidase 4A (*Tm*Pbg4A), assigned the E.C. number 3.2.1.86, is a derivative of *Thermotoga maritima*. It is a 6-phospho-β-glucosidase. The recombinant *Tm*Pbg4A, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 4 (GH4) enzyme (see more details at <a href="https://www.cazy.org">www.cazy.org</a>). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

## **Electrophoretic Purity**

The molecular integrity and purity of *Tm*Pbg4A (GH4) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1**. SDS-PAGE analysis of *Tm*Pbg4A (GH4) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 49,72 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### **Substrate specificity**

 $\textit{Tm} Pbg 4A \text{ (GH4) hydrolyses aryl-phospho-} \beta \text{-glucosides, such as cellobiose 6-phosphate.}$ 

### Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 6.5-10.0 and at a temperature of 37°C. Maximal enzymatic activity is achieved at pH 8 and a consistent temperature of 37°C.

# **Enzyme activity**

The substrate specificity and kinetic properties of *Tm*Pbg4A (GH4) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

#### Reference

Yip et al. (2004) J Am Chem Soc. 126(27):8354-5.

Varrot et al. (2005) J. Mol. Biol. 346:423-435.

#### **Customer Support**

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

### **Quality control assay**

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.