

CZ0312_UG_EN_V2302

Oligoalginate lyase 17A, Saccharophagus degradans

SdAlg17A (PL17)

Catalogue number Presentation

CZ03121 1 mg CZ03122 3 x 1 mg

Description

Oligoalginate lyase 17A (SdAlg17A), assigned the E.C. number 4.2.2.26, is a derivative of Saccharophagus degradans. It is an enzyme that catalysis the degradation of alginate oligosaccharides by a β -elimination mechanism. The recombinant SdAlg17A, purified from Escherichia coli, is a single-domain Pectate Lyase family 17 (PL17) enzyme (see more details at www.cazy.org). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl₂, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

Electrophoretic Purity

The molecular integrity and purity of SdAlg17A (PL17) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).




Figure 1. SDS-PAGE analysis of *Sd*Alg17A (PL17) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 81,36 kDa. Lane M contains a Protein Marker for reference.

Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

Substrate specificity

SdAlg17A (PL17) participates in the eliminative cleavage of low-viscosity alginate.

Temperature and pH optima

The pH optimum for enzymatic activity is 6 while temperature optimum is 40 °C.

Enzyme activity

The substrate specificity and kinetic properties of SdAlg17A (PL17) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

Reference

Kim et al. (2012) Appl Microbiol Biotechnol. 93(5):2233-9.

Park et al. (2014) J Biol Chem. 289(12):8645-55.

Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

Quality control assay

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.