

## Oligosaccharide reducing-end xylanase 8A, *Bifidobacterium adolescentis*

### BaRex8A (GH8)


Catalogue number	Presentation
CZ03851	1 mg
CZ03852	3 x 1 mg

#### Description

Oligosaccharide reducing-end xylanase 8A (*BaRex8A*), assigned the E.C. number 3.2.1.156, is a derivative of *Bifidobacterium adolescentis*. It is an enzyme that participates in the hydrolysis of 1-4- $\beta$ -xylose residues from the reducing end of oligosaccharides. The recombinant *BaRex8A*, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 8 (GH8) enzyme (see more details at [www.cazy.org](http://www.cazy.org)). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

#### Electrophoretic Purity

The molecular integrity and purity of *BaRex8A* (GH8) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1.** SDS-PAGE analysis of *BaRex8A* (GH8) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 45,88 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### Substrate specificity

*BaRex8A* (GH8) hydrolyses wheat flour arabinoxylan and p-nitrophenyl- $\alpha$ -L-arabinofuranoside.

#### Temperature and pH optima

The pH optimum for enzymatic activity is 5 while temperature optimum is 40 °C.

## Enzyme activity

The substrate specificity and kinetic properties of *Ba*Rex8A (GH8) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

## Reference

van den Broek *et al.* (2005) *Appl Microbiol Biotechnol.* 67(5):641-7.

Lagaert *et al.* (2007) *Appl Environ Microbiol.* 73(16):5374-7.

## Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at [info@nzytech.com](mailto:info@nzytech.com).

## Quality control assay

Protein purity is determined to be  $\geq 90\%$ , as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.

---

**NZYtech Lda.** Estrada do Paço do Lumiar, Campus do Lumiar - Edifício E, R/C, 1649-038 Lisboa, Portugal Tel.:+351.213643514 Fax:  
+351.217151168 [www.nzytech.com](http://www.nzytech.com)