

CZ0432 UG EN V2302

Oligoalginate lyase 7A, Pseudomonas aeruginosa

PaAlg7A (PL7)

Catalogue number Presentation

CZ04321 1 mg CZ04322 3 x 1 mg

Description

Oligoalginate lyase 7A (PaAlg7A), assigned the E.C. number 4.2.2.-, is a derivative of $Pseudomonas\ aeruginosa$. It is an enzyme that catalysis the degradation of alginate oligosaccharides by a β -elimination mechanism. The recombinant PaAlg7A, purified from $Escherichia\ coli$, is a single-domain Pectate Lyase family 7 (PL7) enzyme (see more details at www.cazy.org). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl₂, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

Electrophoretic Purity

The molecular integrity and purity of *Pa*Alg7A (PL7) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).




Figure 1. SDS-PAGE analysis of *Pa*Alg7A (PL7) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 27,21 kDa. Lane M contains a Protein Marker for reference.

Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

Substrate specificity

PaAlg7A (PL7) participates in the eliminative cleavage of alginate (preferably poly(MG) block structure).

Temperature and pH optima

The pH optimum for enzymatic activity is 8.5 while temperature optimum is 40 °C.

Enzyme activity

The substrate specificity and kinetic properties of PaAlg7A (PL7) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

Reference

Yamasaki et al. (2003) Acta Crystallogr D Biol Crystallogr. 59(Pt 8):1499-501.

Yamasaki et al. (2004) J. Biol. Chem. 279(30): 31863-31872.

Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

Quality control assay

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.