

CZ0814\_UG\_EN\_V2302

# Pectin acetyl Esterase 12A, Dickeya dadantii

# DdPae12A (CE12)

Catalogue number Presentation


CZ08141 1 mg CZ08142 3 x 1 mg

### **Description**

Pectin acetyl Esterase 12A (*Dd*Pae12A), assigned the E.C. number 3.1.1.-, is a derivative of *Dickeya dadantii*. It is an enzyme that participates in the deacetylation of pectins. The recombinant *Dd*Pae12A, purified from *Escherichia coli*, is a single-domain Carbohydrate Esterase family 12 (CE12) enzyme (see more details at <a href="https://www.cazy.org">www.cazy.org</a>). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

#### **Electrophoretic Purity**

The molecular integrity and purity of *Dd*Pae12A (CE12) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1**. SDS-PAGE analysis of *Dd*Pae12A (CE12) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 60,10 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### **Substrate specificity**

DdPae12A (CE12) participates in the de-esterification of triacetin and sugar beet pectin.

## Temperature and pH optima

The pH optimum for enzymatic activity is 8 while temperature optimum is 30 °C.

# **Enzyme activity**

The substrate specificity and kinetic properties of *Dd*Pae12A (CE12) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

#### Reference

Shevchik and Hugouvieux-Cotte-Pattat. (1997) Mol Microbiol. 24(6):1285-301.

# **Customer Support**

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

# **Quality control assay**

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.