

## $\beta$ -Acetylglucosaminidase 18A, *Enterococcus faecalis*

### EfAcp18A (GH18)


Catalogue number	Presentation
CZ08371	1 mg
CZ08372	3 x 1 mg

#### Description

$\beta$ -Acetylglucosaminidase 18A (EfAcp18A), assigned the E.C. number 3.2.1.96, is a derivative of *Enterococcus faecalis*. It is an enzyme that participates in the endohydrolysis of the diacetylchitobiosyl unit in high-mannose glycopeptides and glycoproteins containing the (Man(GlcNAc)(2))Asn-structure: one N-acetyl-D-glucosamine residue remains attached to the protein, the rest of the oligosaccharide is released intact. The recombinant EfAcp18A, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 18 (GH18) enzyme (see more details at [www.cazy.org](http://www.cazy.org)). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

#### Electrophoretic Purity

The molecular integrity and purity of EfAcp18A (GH18) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1.** SDS-PAGE analysis of EfAcp18A (GH18) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 32,22 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### Substrate specificity

EfAcp18A (GH18) hydrolyses N-linked glycans from glycoproteins.

## Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 6.0-7.0 and at a temperature of 37°C. Maximal enzymatic activity is achieved at pH 6.5 and a consistent temperature of 37°C.

## Enzyme activity

The substrate specificity and kinetic properties of *EfAcp18A* (GH18) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

## Reference

Bøhle *et al.* (2011) FEMS Microbiol Lett. 325(2):123-9.

## Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at [info@nzytech.com](mailto:info@nzytech.com).

## Quality control assay

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.

---

**NZYtech Lda.** Estrada do Paço do Lumiar, Campus do Lumiar - Edifício E, R/C, 1649-038 Lisboa, Portugal Tel.: +351.213643514 Fax: +351.217151168 [www.nzytech.com](http://www.nzytech.com)