

CZ0848\_UG\_EN\_V2302

# Rhamnogalacturonan thetaiotaomicron

### rhamnohydrolase

106A,

**Bacteroides** 

## BtRhh106A (GH106)


Catalogue numberPresentationCZ084811 mgCZ084823 x 1 mg

#### Description

Rhamnogalacturonan rhamnohydrolase 106A (BtRhh106A), assigned the E.C. number 3.2.1.174, is a derivative of Bacteroides thetaiotaomicron. It is an enzyme that participates in the exohydrolysis of the  $\alpha$ -L-Rha-1,4- $\alpha$ -D-GalA bond in rhamnogalacturonan oligosaccharides with initial inversion of configuration releasing  $\beta$ -L-rhamnose from the non-reducing end of rhamnogalacturonan oligosaccharides. The recombinant BtRhh106A, purified from Escherichia coli, is a single-domain Glycoside Hydrolase family 106 (GH106) enzyme (see more details at <a href="https://www.cazy.org">www.cazy.org</a>). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

#### **Electrophoretic Purity**

The molecular integrity and purity of *Bt*Rhh106A (GH106) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1**. SDS-PAGE analysis of *Bt*Rhh106A (GH106) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 104,8 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### **Substrate specificity**

 $\it Bt$ Rhh106A (GH106) hydrolyses the L-Rha-lpha4-D-GalA linkage in rhamnogalacturonan I (RGI).

#### Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 6.5-7.5 and at a temperature of 37°C. Maximal enzymatic activity is achieved at pH 7 and a consistent temperature of 37°C.

#### **Enzyme activity**

The substrate specificity and kinetic properties of *Bt*Rhh106A (GH106) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

#### Reference

Luis et al. (2018) Nat Microbiol. 3(2):210-219.

#### **Customer Support**

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

#### **Quality control assay**

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.