

CZ0851 UG EN V2302

## **β-Galactosidase 42A**, Bacteroides thetaiotaomicron

# BtLac43A-Lac42A (GH43-GH42)

Catalogue number Presentation


CZ08511 1 mg CZ08512 3 x 1 mg

#### **Description**

β-Galactosidase 42A (*Bt*Lac43A-Lac42A), assigned the E.C. number 3.2.1.23, is a derivative of *Bacteroides thetaiotaomicron*. It is a β-galactosidase. The recombinant *Bt*Lac43A-Lac42A, purified from *Escherichia coli*, is a modular Glycoside Hydrolase family 42 (GH43-GH42) enzyme (see more details at <a href="https://www.cazy.org">www.cazy.org</a>). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

#### **Electrophoretic Purity**

The molecular integrity and purity of *Bt*Lac43A-Lac42A (GH43-GH42) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1**. SDS-PAGE analysis of *Bt*Lac43A-Lac42A (GH43-GH42) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 113,8 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### **Substrate specificity**

BtLac43A-Lac42A (GH43-GH42) hydrolyses rhamnogalacturonan I (RGI).

#### Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 6.5-7.5 and at a temperature of 37°C. Maximal enzymatic activity is achieved at pH 7 and a consistent temperature of 37°C.

## **Enzyme activity**

The substrate specificity and kinetic properties of BtLac43A-Lac42A (GH43-GH42) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

#### Reference

Mahowald et al. (2009) Proc Natl Acad Sci U S A. 106(14):5859-5864.

## **Customer Support**

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

## **Quality control assay**

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.