

# Unsaturated rhamnogalacturonyl hydrolase 105C, *Bacteroides thetaiotaomicron*

## *BtUrh105C* (GH105)


Catalogue number	Presentation
CZ08831	1 mg
CZ08832	3 x 1 mg

### Description

Unsaturated rhamnogalacturonyl hydrolase 105C (*BtUrh105C*), assigned the E.C. number 3.2.1.-, is a derivative of *Bacteroides thetaiotaomicron*. It catalyzes the hydrolysis of unsaturated rhamnogalacturonan II backbone. The recombinant *BtUrh105C*, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 105 (GH105) enzyme (see more details at [www.cazy.org](http://www.cazy.org)). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

### Electrophoretic Purity

The molecular integrity and purity of *BtUrh105C* (GH105) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1.** SDS-PAGE analysis of *BtUrh105C* (GH105) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 45,76 kDa. Lane M contains a Protein Marker for reference.

### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

### Substrate specificity

*BtUrh105C* (GH105) hydrolyses the D-Gal- $\alpha$ 1,4-D-Gal linkage in pectins in particular of rhamnogalacturonan II (RGII).

## Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 6.5-7.5 and at a temperature of 37°C. Maximal enzymatic activity is achieved at pH 7 and a consistent temperature of 37°C.

## Enzyme activity

The substrate specificity and kinetic properties of *BtUrh105C* (GH105) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

## Reference

Ndeh *et al.* (2017) *Nature*. 544(7648):65-70.

Mahowald *et al.* (2009) *Proc Natl Acad Sci U S A*. 106(14):5859-64.

## Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at [info@nzytech.com](mailto:info@nzytech.com).

## Quality control assay

Protein purity is determined to be  $\geq 90\%$ , as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.

---

**NZYtech Lda.** Estrada do Paço do Lumiar, Campus do Lumiar - Edifício E, R/C, 1649-038 Lisboa, Portugal Tel.:+351.213643514 Fax:  
+351.217151168 [www.nzytech.com](http://www.nzytech.com)