

CZ0940_UG_EN_V2302

Alginate lyase 7A, Vibrio sp.

VAlg7A (PL7)

Catalogue numberPresentationCZ094010.5 mgCZ094023 x 0.5 mg

Description

Alginate lyase 7A (VAlg7A), assigned the E.C. number 4.2.2.3, is a derivative of Vibrio sp.. It is an enzyme that catalysis the degradation of alginate by a β -elimination mechanism. The recombinant VAlg7A, purified from E scherichia coli, is a single-domain Pectate Lyase family 7 (PL7) enzyme (see more details at $\underline{www.cazy.org}$). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl₂, and 25% (v/v) glycerol, at a concentration of 0.5 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

Electrophoretic Purity

The molecular integrity and purity of VAlg7A (PL7) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).




Figure 1. SDS-PAGE analysis of VAlg7A (PL7) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 31,73 kDa. Lane M contains a Protein Marker for reference.

Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

Substrate specificity

VAlg7A (PL7) participates in the eliminative cleavage of sodium alginate.

Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 7.6-9.0 and at a temperature of 30°C. Maximal enzymatic activity is achieved at pH 8.5 and a consistent temperature of 30°C.

Enzyme activity

The substrate specificity and kinetic properties of VAlg7A (PL7) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

Reference

Uchimura et al. (2010) Mar Biotechnol (NY) 12(5):526-533.

Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

Quality control assay

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.