

CZ0945_UG_EN_V2302

Xylanase 98A, Ruminococcus champanellensis

RcXyn98A (GH98-CBM35)

Catalogue numberPresentationCZ094510.5 mgCZ094523 x 0.5 mg

Description

Xylanase 98A (*Rc*Xyn98A), assigned the E.C. number 3.2.1.8, is a derivative of *Ruminococcus champanellensis*. It is an endo-1,4-β-xylanase. The recombinant *Rc*Xyn98A, purified from *Escherichia coli*, is a modular Glycoside Hydrolase family 98 (GH98-CBM35) enzyme (see more details at www.cazy.org). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl₂, and 25% (v/v) glycerol, at a concentration of 0.5 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

Electrophoretic Purity

The molecular integrity and purity of *Rc*Xyn98A (GH98-CBM35) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).




Figure 1. SDS-PAGE analysis of *Rc*Xyn98A (GH98-CBM35) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 113,1 kDa. Lane M contains a Protein Marker for reference.

Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

Substrate specificity

RcXyn98A (GH98-CBM35) hydrolyses xylans in particular beechwood xylan.

Temperature and pH optima

The pH optimum for enzymatic activity is 6 while temperature optimum is 37 °C.

Enzyme activity

The substrate specificity and kinetic properties of *Rc*Xyn98A (GH98-CBM35) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

Reference

Moraïs et al. (2016) Environ Microbiol. 18(2):542-56.

Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

Quality control assay

Protein purity is determined to be ≥75%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.