

CZ0946\_UG\_EN\_V2302

## Cellulase 9B, Thermobifida fusca

# TfCel9B (CBM4-CBM30-GH9-CBM2)


Catalogue number	Presentation
CZ09461	1 mg
CZ09462	3 x 1 mg

## Description

Cellulase 9B (*Tf*Cel9B), assigned the E.C. number 3.2.1.4, is a derivative of *Thermobifida fusca*. It is an endo-1,4- $\beta$ -glucanase. The recombinant *Tf*Cel9B, purified from *Escherichia coli*, is a modular Glycoside Hydrolase family 9 (CBM4-CBM30-GH9-CBM2) enzyme (see more details at <u>www.cazy.org</u>). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

#### **Electrophoretic Purity**

The molecular integrity and purity of *Tf*Cel9B (CBM4-CBM30-GH9-CBM2) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1**. SDS-PAGE analysis of *Tf*Cel9B (CBM4-CBM30-GH9-CBM2) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 102,4 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### Substrate specificity

*Tf*Cel9B (CBM4-CBM30-GH9-CBM2) hydrolyses CMC, swollen cellulose and pNP-cellobiose.

#### Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 5.5-6.0 and at a temperature of 50°C. Maximal enzymatic activity is achieved at pH 5.5 and a consistent temperature of 50°C.

## **Enzyme activity**

The substrate specificity and kinetic properties of *Tf*Cel9B (CBM4-CBM30-GH9-CBM2) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

## Reference

Irwin et al. (1993) Biotechnol Bioeng. 42(8):1002-13.

Sakon et al. (1997) Nat Struct Biol. 4(10):810-8.

#### **Customer Support**

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

#### **Quality control assay**

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.