

## Endo- $\alpha$ -N-acetylgalactosaminidase 101A, *Clostridium perfringens*

### CpNga101A (GH101)


Catalogue number	Presentation
CZ09551	0.5 mg
CZ09552	3 x 0.5 mg

#### Description

Endo- $\alpha$ -N-acetylgalactosaminidase 101A (CpNga101A), assigned the E.C. number 3.2.1.97, is a derivative of *Clostridium perfringens*. It is an enzyme that catalyzes the liberation of Gal-1,3- $\beta$ -GalNAc  $\alpha$ -linked to serine or threonine residues of mucin-type glycoproteins. The recombinant CpNga101A, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 101 (GH101) enzyme (see more details at [www.cazy.org](http://www.cazy.org)). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 0.5 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

#### Electrophoretic Purity

The molecular integrity and purity of CpNga101A (GH101) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1.** SDS-PAGE analysis of CpNga101A (GH101) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 128,2 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### Substrate specificity

CpNga101A (GH101) hydrolyses serine/threonine glycoproteins.

#### Temperature and pH optima

The pH optimum for enzymatic activity is 6 while temperature optimum is 60 °C.

## Enzyme activity

The substrate specificity and kinetic properties of CpNga101A (GH101) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

## Reference

Ashida *et al.* (2008) *Glycobiology*. 18(9):727-34.

## Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at [info@nzytech.com](mailto:info@nzytech.com).

## Quality control assay

Protein purity is determined to be  $\geq 75\%$ , as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.

---

**NZYtech Lda.** Estrada do Paço do Lumiar, Campus do Lumiar - Edifício E, R/C, 1649-038 Lisboa, Portugal Tel.:+351.213643514 Fax:  
+351.217151168 [www.nzytech.com](http://www.nzytech.com)