

CZ0984 UG EN V2302

# Cellulase 9A, Clostridium papyrosolvens

# CpCel9A (GH9-CBM3-doc-doc)


Catalogue numberPresentationCZ098410.5 mgCZ098423 x 0.5 mg

### **Description**

Cellulase 9A (CpCel9A), assigned the E.C. number 3.2.1.4, is a derivative of  $Clostridium\ papyrosolvens$ . It is an endo-1,4- $\beta$ -glucanase. The recombinant CpCel9A, purified from  $Escherichia\ coli$ , is a modular Glycoside Hydrolase family 9 (GH9-CBM3-doc-doc) enzyme (see more details at  $\underline{www.cazy.org}$ ). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 0.5 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

#### **Electrophoretic Purity**

The molecular integrity and purity of *Cp*Cel9A (GH9-CBM3-doc-doc) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1**. SDS-PAGE analysis of *Cp*Cel9A (GH9-CBM3-doc-doc) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 76,99 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### **Substrate specificity**

CpCel9A (GH9-CBM3-doc-doc) hydrolyses CMC and avicel.

### Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 6.5-7.5 and at a temperature of 37°C. Maximal enzymatic activity is achieved at pH 7 and a consistent temperature of 37°C.

# **Enzyme activity**

The substrate specificity and kinetic properties of *Cp*Cel9A (GH9-CBM3-doc-doc) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

#### Reference

Zepeda et al. (2013) Genome Announc. 1(5): e00698-13.

# **Customer Support**

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

# **Quality control assay**

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.