

CZ0989 UG EN V2302

# Fucosidase 95A, Clostridium perfringens

# CpFuc95A (GH95)

Catalogue number Presentation


CZ09891 1 mg CZ09892 3 x 1 mg

#### **Description**

Fucosidase 95A (CpFuc95A), assigned the E.C. number 3.2.1.63, is a derivative of *Clostridium perfringens*. It is a 1,2- $\alpha$ -L-fucosidase. The recombinant CpFuc95A, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 95 (GH95) enzyme (see more details at <a href="https://www.cazy.org">www.cazy.org</a>). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

### **Electrophoretic Purity**

The molecular integrity and purity of *Cp*Fuc95A (GH95) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1**. SDS-PAGE analysis of *Cp*Fuc95A (GH95) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 91,99 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### **Substrate specificity**

CpFuc95A (GH95) hydrolyses fucosyl glycans and porcine gastric mucin.

#### Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 4.0-11.0 and at a temperature of 60°C. Maximal enzymatic activity is achieved at pH 8 and a consistent temperature of 60°C.

# **Enzyme activity**

The substrate specificity and kinetic properties of *Cp*Fuc95A (GH95) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

#### Reference

Fan et al. (2016) J Basic Microbiol. 56(4):347-57.

# **Customer Support**

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

# **Quality control assay**

Protein purity is determined to be ≥75%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.