

## $\beta$ -L-Arabinofuranosidase 127A, *Bacteroides thetaiotaomicron*

### *BtHyp127B* (GH127)


Catalogue number	Presentation
CZ10033	1 mg
CZ10034	3 x 1 mg

#### Description

$\beta$ -L-Arabinofuranosidase 127A (*BtHyp127B*), assigned the E.C. number 3.2.1.185, is a derivative of *Bacteroides thetaiotaomicron*. It is a  $\beta$ -L-arabinofuranosidase. The recombinant *BtHyp127B*, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 127 (GH127) enzyme (see more details at [www.cazy.org](http://www.cazy.org)). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

#### Electrophoretic Purity

The molecular integrity and purity of *BtHyp127B* (GH127) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1.** SDS-PAGE analysis of *BtHyp127B* (GH127) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 76,91 kDa. Lane M contains a Protein Marker for reference.

#### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

#### Substrate specificity

*BtHyp127B* (GH127) hydrolyses the L-Acef- $\alpha$ 1,3-L-Rhap linkage in rhamnogalacturonan II (RGII).

#### Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 6.5-7.5 and at a temperature of 37°C. Maximal enzymatic activity is achieved at pH 7 and a consistent temperature of 37°C.

## Enzyme activity

The substrate specificity and kinetic properties of *BtHyp127B* (GH127) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

## Reference

Cartmell *et al.* (2018) *Nat Microbiol.* 3(11):1314-1326.

Mahowald *et al.* (2009) *Proc Natl Acad Sci U S A.* 106(14):5859-64.

## Customer Support

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at [info@nzytech.com](mailto:info@nzytech.com).

## Quality control assay

Protein purity is determined to be  $\geq 90\%$ , as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.