

CZ1024\_UG\_EN\_V2302

# β-L-Arabinofuranosidase 142A, Bacteroides thetaiotaomicron

# *Bt*Hyp142A (GH142)


Catalogue number	Presentation
CZ10241	1 mg
CZ10242	3 x 1 mg

### Description

 $\beta$ -L-Arabinofuranosidase 142A (*Bt*Hyp142A), assigned the E.C. number 3.2.1.185, is a derivative of *Bacteroides thetaiotaomicron*. It is a  $\beta$ -L-arabinofuranosidase. The recombinant *Bt*Hyp142A, purified from *Escherichia coli*, is a single-domain Glycoside Hydrolase family 142 (GH142) enzyme (see more details at <u>www.cazy.org</u>). The protein is supplied in a solution containing 35 mM NaHepes buffer (pH 7.5), 750 mM NaCl, 200 mM Imidazole, 3.5 mM CaCl<sub>2</sub>, and 25% (v/v) glycerol, at a concentration of 1 mg/mL. Bulk quantities of this product can be made available upon request. To place an order, simply visit our website. We offer fast and secure shipping worldwide.

### **Electrophoretic Purity**

The molecular integrity and purity of *Bt*Hyp142A (GH142) were evaluated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), followed by BlueSafe staining (MB15201) (Figure 1).



**Figure 1**. SDS-PAGE analysis of *Bt*Hyp142A (GH142) was conducted in (Lane 1), employing a 14% polyacrylamide gel. The enzyme exhibits a band corresponding to a molecular weight of approximately 60,22 kDa. Lane M contains a Protein Marker for reference.

### Storage temperature

The protein should be stored at -30°C to -15°C in a constant temperature freezer. The protein will remain stable till the expiry date if stored as specified.

### Substrate specificity

BtHyp142A (GH142) hydrolyses arabinan in chain D of rhamnogalacturonan II (RGII).

# Temperature and pH optima

The enzyme exhibits optimal activity within a pH range of 6.5-7.5 and at a temperature of 37°C. Maximal enzymatic activity is achieved at pH 7 and a consistent temperature of 37°C.

# **Enzyme activity**

The substrate specificity and kinetic properties of *Bt*Hyp142A (GH142) are detailed in the referenced publication provided below. To perform enzyme assays and determine specific activity values, adhere to the methodology outlined in the cited paper(s).

### Reference

Ndeh et al. (2017) Nature. 544(7648):65-70.

# **Customer Support**

Our dedicated customer support team is always ready to assist you with any questions or technical issues you may have. Reach us via email at info@nzytech.com.

# **Quality control assay**

Protein purity is determined to be ≥90%, as assessed by SDS-PAGE and subsequent BlueSafe staining (MB15201).

For life science research only. Not for use in diagnostic procedures.

NZYtech Lda. Estrada do Paço do Lumiar, Campus do Lumiar - Edifício E, R/C, 1649-038 Lisboa, Portugal Tel.:+351.213643514 Fax: +351.217151168 www.nzytech.com